

ENHANCEMENT OF IMMUNITY, ANTIOXIDANT STATUS, METABOLIC ATTRIBUTES AND REPRODUCTIVE HORMONAL PROFILES BY ADDING SOME NATURAL ADDITIVES TO RABBIT DOE'S DIET

M.E. El-Speiy¹, M.A. El-Sawy¹, T.A. Sadak¹, B.M. Abou-Shehema¹, H.A. Shahba¹, M.R. Habib¹, M.M. Abdella¹, A.M. Tammam², and M.M. Zeitoun^{3*}

¹Animal Production Research Institute, Agriculture Research Center, Egypt.

²Poultry Nutrition Department, Faculty of Agriculture, Ain Shams University, Egypt.

³Department of Animal and Fish Production, Faculty of Agriculture, Alexandria University, Egypt.

e. mail: Elsawy1966 @ gmail. com, Doctor.tarek2@ gmail.com, Mrh20152020@ gmail.com, abdellahmm@ yahoo.com, mmzeitoun@yahoo.com

*Corresponding author: mohamed.elspeiy@yahoo.com

ABSTRACT

The aim of the study aimed to find out cheap local bioactive materials that support immunity, antioxidant capacity, and reproductive efficiency of California rabbits. Seventy-two adult rabbit does were equally allocated into six treatments (12, each). All does were fed a basal diet and orally given three ml distilled water as a basal solution during 10 days before mating until 28-days post insemination. Control does (C) received no supplement, G2 (HB) does were given 50mg honeybee/head/day; G3 (DPP) received 50 mg date palm pollen/head/day; G4 (BP) received 200mg bee pollen/head/day; G5 (SY) received three ml of a commercial synbiotic, and G6 (MIX) received a combination of all previous compounds.

Results indicated that all treatments improved ($P<0.05$) the immunity

functions and antioxidant biomarkers. Similarly, all treatments increased ($P<0.05$) plasma protein, albumin, and globulins, However it reduced ($P<0.05$) liver enzymes activity. Treatments remarkably reduced the bad cholesterol (free and LDL) and triglycerides, but increased HDL compared with control. Furthermore, all treatments elicited more secretion of ovarian hormones, which in turn increased pituitary gonadotrophins (FSH and LH). Contrariwise, all treatments resulted in lower levels of prolactin.

In conclusion, provision of rabbit does during its productive life with bee products, date palm pollen and synbiotic would be a good solution for improving productive and reproductive performance of such animal.

Keywords: Bee products, date palm, immunity, probiotic, productivity,

INTRODUCTION

The nutritional composition of some medicinal plants is believed to benefit productive, reproductive and performance in mammals; specifically, it has been found that these nutritive compounds are able to treat sexual dysfunction (**Sumalatha *et al*, 2010**). Moreover, the consumers awareness of food safety is expanding rapidly. Public health concerns are emerged over the use of antibiotics as growth promoters in farm animals' feed (**De Marco *et al*, 2015**). Therefore, great concern has been raised to find out natural alternatives for animal feeding strategies (**Hosseini *et al*, 2013**). As a result of the emerging information, the feed industry began searching for alternative feed additives to replace antibiotics in order to provide proper productivity. Antibiotics and other pharmaceuticals have long been widely utilized, with the objective of manipulating gut microbes for enhancing food for animal productivity. Long-term use of some of these drugs has led to the formation of drug-resistant microorganisms, representing a health threat to the consumers leading to a negative impact on the human gut ecology (**Biernasiak *et al*, 2010**).

Synbiotics (SY) are essential ingredients in animal feeding because they include both probiotics and prebiotics (**Gibson and Roberfroid, 1995**). Synbiotic improves host welfare by enhancing the survival and activity of live microbial dietary supplements in the digestive tract, by selectively stimulating the development and/or activating the metabolism of one specie or more of health-promoting bacteria (**Cencic and Chingwaru, 2010**).

Bee pollen (BP) could also be a promising additive since it involves a significant number of bioactive compounds such as carbohydrates, enzymes, vitamins, fatty acids, essential amino acids or carotenoids depending on the botanical and geographical origin of the bee pollen (**Denisow and Denisow-Pietrzyk 2016**). Additionally, bee pollen contains bioactive components including; polyphenolics, flavonoids, proteins, and essential amino acids (**Carpes *et al*, 2009**). Furthermore, BP contains vitamins, minerals and coenzymes which are recognized for their antioxidant activities and contribute to anti-inflammatory effects (**Attia *et al*, 2011 and Morais *et al*, 2011**). The variability of metabolites in bee pollen sets this product apart from other bee products and can be used in a variety of medical and therapeutic applications. Date palm pollen (DPP) is also known to include variety of compounds, including high total phenolics content, flavonoids, anthocyanins, and seleno-proteins (**Baliga *et al*, 2011**), making it an excellent candidate for antioxidant activity with little negative side effects (**Fallahi *et al*, 2015**). Moreover,

honeybee (HB) contains a high amount of metabolizable energy in the form of glucose and fructose, and also antibacterial activity against microorganisms that inhibit a broad spectrum of bacteria, including aerobes and anaerobes, gram positives and negatives (**Hannan et al, 2009**). Seemingly, HB contains low levels of amino acids and vitamins, antioxidant properties, phenolic acids and flavonoids (**Andrade et al, 1997**). However, it includes enzymes, ascorbic acid, and minerals (**White, 1975**). Thus, honeybees have antimicrobial properties, according to **Molan and Russell (1988)**. Therefore, the use of natural products as dietary supplements could be a reasonable alternative for enhancing immunity, antioxidant status, metabolic, and hormonal profiles of rabbits raised under shortage of feedstuffs in hot-arid developing countries.

Therefore, the aim of the present study to find out cheap some local bioactive materials that support immunity, antioxidant capacity, and reproductive efficiency of California rabbits.

MATERIALS AND METHODS

The study was conducted at a private rabbit farm in Qalyubia governorate, where ambient indoor temperatures ranged from 19 to 27°C, humidity 43 to 54 %, and the light/dark cycle was 16/8 hours throughout the study term.

Experimental design:

In this study, seventy-two Californian does aged 6 months, and weighted of 3250 ± 78.2 g were used. Rabbits were randomly allotted into 6 equal treatment groups (12 in each), and orally received 3 ml distilled water daily for 10 days before mating until 28 days post mating. Control (C) does received distilled water only, however treated does received the tested bioactive compounds.

All does were fed a basal diet (Table 1), and submitted to the following treatments; control (C) received no supplement, Synbiotic does (SY) received 3ml synbiotic (0.1%), Date palm pollen(DPP) does received 50mg per a doe/day, Bee pollen(BP) does were given 200mg per a doe/day, Honeybee (HB) was supplemented with 50 mg per a doe/day, and Mixture (MIX) does received 3 ml synbiotic containing 50 mg DPP 0.200 mg BP, and 50 mg HB. The does were artificially inseminated. Insemination was performed with semen collected and pooled from 15 adult Californian males. The chosen

doses of bee pollen (BP) and date palm pollen (DPP) were according to **El-Hanoun *et al* (2007)**, while synbiotic dose was according to **Abdel-Raheem *et al* (2012)**.

Housing, feeding and management:

Rabbits were housed in a naturally ventilated barn equipped with wire galvanized batteries (60 × 55 × 40 cm, L×W×D). The batteries were equipped with pellet feeders and automatic drinkers. Fresh water was provided in the experiment. The basal experimental diet (Table 1) was designed and pelleted to satisfy the nutritional requirements of rabbits as a whole (**NRC, 1977**).

Blood sampling:

Blood samples (5ml) were withdrawn at the end of the trial from marginal ear veins from all does within a treatment in the morning before receiving feed and water, using sterile disposable needles and heparinized sterile tubes. Plasma samples were obtained after centrifugation blood samples at 3000 rpm for 15 min and stored at -20°C until analyzed.

Hormonal determinations:

Plasma progesterone (P4) was determined according to the method of **Simersky *et al* (2007)** using a commercial ELISA 96-well kit (Diagnostics Test Canada, Inc., Ontario, Canada). The intra-assay coefficient of variation (C.V) was 7.6%.

Plasma estradiol-17β (E2) was analyzed according to the method of **Ratcliff *et al* (1988)** using enzyme-linked immune absorbent assay (ELISA) kits (Diagnostics Test Canada, Inc., Ontario, Canada). The intra-assay C.V was 7.2%. All samples were analyzed in one assay for each hormone.

Plasma prolactin (PRL) determinations were done according to **Zoli *et al* (2002)** using a commercial ELISA kit (Immuno spec Corporation/ USA). The intra-assay CV was 8.3%.

Plasma FSH determination was done by a commercial ELISA kit (My Bio Source, San Diego, CA, USA). The intra-assay CV was 7.8%. Plasma LH values was determined by a commercial ELISA kit (My BioSource, San Diego, CA, USA). The intra-assay CV was 7.6%.

Blood biochemical determinations:

Alanine aminotransferase (ALT) and aspartate aminotransferase (AST) activities were accomplished according to **Reitman and Frankel (1957)**. Total protein (TP) was determined according to **Song *et al* (2023)**, albumin (Alb) was

Table 1. Experimental diet composition and chemical analysis.

Ingredient	%	Calculated analysis	
Yellow corn	6.22	Crude protein, %	18.8
Soybean meal, 44%	22.33	Crude fiber, %	13.0
Wheat bran	23.33	Ether extract, %	3.0
Barley	15.00	Digestible energy (kcal/kg diet)	2680
Alfalfa hay	30.12	n-6 poly unsaturated FAs%	0.3
Ground limestone	1.00	n-3 poly unsaturated FAs%	1.03
Dicalcium phosphate	1.20	Determined analysis (g/kg)	
Common salt	0.50	Dry matter	897.1
Vit. + min. premix*	0.30	Organic matter	801.4
Total	100.0	Crude protein	169.8
		Crude fiber	138.5
		Ether extract	26.2
		Nitrogen-free extract	575.0
		Ash	87.9

*Each 3 kg of premix contains: Vit. A: 12, 000,000 IU; Vit. D₃: 3, 000,000 IU; Vit. E: 10.0 mg; Vit. K₃: 3.0 mg; Vit. B₁: 200 mg; Vit. B₂: 5.0 mg; Vit. B₆: 3.0 mg; Vit. B₁₂: 15.0 mg; Biotin: 50.0 mg; Folic acid: 1.0 mg; Nicotinic acid: 35.0 mg; Pantothenic acid: 10.0 mg; Mn: 80 g; Cu: 8.8 g; Zn: 70 g; Fe: 35 g; I: 1 g; Co: 0.15g and Se: 0.3g.

quantified according to **Doumas et al (1971)**, however, globulin (Glo) was estimated by the difference between TP and Alb. Triglycerides (TG) levels were determined using the method of **Fossati and Prencipe (1982)**. Determinations of total cholesterol (TC), high-density lipoprotein (HDL) and low-density lipoprotein (LDL) were done following the method of **Birtcher and Ballantyne (2004)**. All biochemical attributes were determined in commercial kits (Human, Liquicolor, Germany).

Antioxidant activity determination:

Plasma total antioxidant capacity (TAC) and malondialdehyde (MDA) were determined using commercially available kits using a spectrophotometer (GNW-Model: SM-721) according to the methods reported by **Gawel et al (2004)**.

Immunoglobulins determination

Rabbit IgG and IgM were determined by commercial ELISA kits (Abcam Limited, Cambridge, UK). The sensitivity of the IgG assay was 0.23ng/mL (**Lopez and Alvariño, 2000**) and the intra-assay CV was 6.5%,

however, the sensitivity of the IgM assay was 0.188 ng/ml, and the intra-assay CV was 7.2%.

Measurement of plasma digestive enzyme:

Amylase and lipase activities were determined using commercial kits (Human Gesellschaft, Germany) by the use of an automated analyzer (Chem-Well 2900T, USA).

Statistical analysis:

All data were subjected to the least square analysis of variance according to **Steel and Torrie (1984)** using SAS package (**SAS, 2002**).

The significant differences among treatment groups were tested using Multiple Range Test according to **Duncan (1955)**.

RESULTS AND DISCUSSION

Immunoglobulins and antioxidants

As shown in Table 2, all bioactive components and their mixture enhanced ($P < 0.05$) both fractions of Ig. The improvement ranges between 44-90% increases above control. Moreover, TAC as an indicator of the anti-oxidation biomarker, remarkably ($P < 0.05$) increased due to treatments. The improvement ranges between 61-116% above control values. Basically, the increase of MDA is a strong indicator of the accumulation of reactive oxygen species. Treatment reduced the levels of MDA by 22-33% of the control value. It has been recently established that bee derivatives (i.e. honey, pollens) are crucial components for increasing the immunity and the anti-oxidative status via improvement of intestinal absorption of nutrients and activating digestive enzymes in large animals, poultry and rabbits (**Abdelnour *et al*, 2019** and **Abd El-Aziz *et al*, 2023**).

The richness of honeybee and bee pollens in simple sugars, vitamins, carotenoids, essential amino acids, polyunsaturated fatty acids, proteins and minerals imposes vital roles for such compounds. The immune-competence requires special nutrients such as bee products (honeybee, royal jelly, bee pollens, venom).

Table 2. Effect of natural additive on blood plasma immune-globulins and antioxidant indicators of Californian rabbit does

Treatment groups	IgG (mg/dl)	IgM (mg/dl)	TAC (nmol/ml)	MDA (nmol/ml)
Control (C)	461.0 ^d	17.0 ^c	1.04 ^c	7.15 ^a
Honeybee (HB)	878.0 ^a	23.0 ^{bc}	2.25 ^a	4.77 ^c
Date Palm Pollen (DPP)	663.0 ^c	28.0 ^b	1.89 ^{ab}	5.49 ^b
Bee pollen (BP)	723.0 ^b	32.0 ^a	1.68 ^b	5.32 ^b
Synbiotic (SY)	727.0 ^b	26.0 ^{ab}	1.87 ^b	4.93 ^c
Mixture (MIX)	858.0 ^a	21.0 ^{bc}	1.79 ^{ab}	5.53 ^b
MSE	8.73	3.49	0.35	0.56
P-value	0.0001	0.0004	0.0001	0.006

a,b,c,d Values with different superscripts in the same column differ significantly ($P < 0.05$). IgG: immunoglobulin G, IgM: Immunoglobulin M, TAC: Total antioxidant capacity, MDA: malondialdehyde.

Blood biochemical changes:

As shown in Table 3 illustrates significant ($P < 0.05$) increases in TP, Alb and Glo. Contrariwise, liver enzymes (AST and ALT) decreased ($P < 0.05$) in does that were given the bioactive compounds, especially when a synbiotic was given. The highest values of protein and its fractions accompanied with the lowest liver enzymes were observed in case of synbiotic treatment. Generally, probiotics enhance the gut health leading to increased absorptive capability of the intestinal mucosa to absorb more protein (**Jäger et al, 2018**). In conjunction with what mentioned in Table 2, protein and its fraction, globulins play vital roles in the immunity system health and function. Albumin regulates the distribution of the extracellular fluids and acts as a carrier for a variety of bio-chemicals such as bilirubin, fatty acids, hormones and vitamins (**Attia et al, 2015**). Additionally, albumin-based antibodies are the main protein components that are synthesized in the hepatic tissues functioning as humoral immune agents. Other research study on sheep concluded significant increases of protein and globulins in lambs provided with probiotic, prebiotic, and synbiotic (**Ellithy et al, 2022**). Furthermore, synbiotic, honeybee, bee pollens, and date palm pollens reduced AST and ALT which means improvement of the liver health. Evidently, the content of flavonoids in date palm pollen are high enough to protect unsaturated fatty acids which strengthen cellular membranes and protect it from oxidation, thus maintain liver integrity and function (**Abdel-Shaheed et al, 2021**). Besides, bee pollen was effective to reduce serum urea, AST and ALT when added to the rabbit'

diet leading to the improvements of hepatic and renal functions (**S'aric' et al, 2009**). This enhancement was attributed to the richness of bee pollen with antioxidant constituents. In the same manner, **El-Hammady et al (2017)** reported increased serum protein and albumin in rabbit bucks provided with bee pollen.

Honeybee is not only a rich source of simple sugars [i.e. glucose (32%) and fructose (43%)], but also it is rich in vitamins, flavonoids, antibacterial and antioxidant properties (**Tuksitha et al, 2018**). Parallel to the present study findings, **El Rabey et al (2013)** revealed that providing male albino rats with 2.5 g/kg body weight natural honeybees significantly improved liver function by reducing AST and ALT activities, and increased serum protein.

Concerning lipid fractions (Table 4), all treatments improved lipid fractions (i.e., reduced LDL and triglycerides and elevated HDL). Only honeybees reduced total cholesterol, however other ingredients tended to decrease total cholesterol. A proposed mechanism of the role of probiotic bacteria on lowering lipid fractions in circulation was stated by **Fukushima and Nakano(1995)** who hypothesized that probiotic bacteria (i.e. *Lactobacilli*, *Bifidobacteria*) pose bile salts hydrolase activity leading to the linkage of cholesterol to the bacteria to produce short-chain fatty acids. Another possible explanation of the reduction of cholesterol and its derivatives is that probiotic microorganisms may inhibit hydroxymethyl-glutaryl co-enzyme A, an enzyme involved in the synthesis of cholesterol (**Abdelhady and El-Abasy, 2015**). Most recently **Kim and Kim (2023)** attributed the cholesterol-lowering effect of dead probiotic bacteria (parabiotic) to three mechanisms; first, binding of probiotic to the dietary cholesterol contributing to its absorption inhibition and promoting its excretion. Second, activating nuclear receptors of enterocytes which modulate cholesterol transport by inhibiting the expression of an influx protein called Niemann-Pick C1-like 1 (NPC1L1) and promoting the expression of an efflux transporter named ATP-Binding Cassette (ABC). Third, reducing the synthesis of lipids in the liver. **Zeedan et al (2017)** stated serum total lipids, cholesterol, and triglycerides were diminished in rabbit bucks given different doses of bee pollen. Clearly, the presence of high levels of unsaturated fatty acids (i.e., oleic, linoleic, and linolenic) could play a vital role in preventing the formation of lipid peroxidation, resulting in reduced levels of triglycerides and cholesterol (**Farag and El-Rayes, 2016**). Additionally, the hypo-lipidemic effect of date palm pollen may be due to presence of polyunsaturated fatty acids and phyto-sterols which interfere with cholesterol absorption via enterocytes (**Abbas and Ateya, 2011**).

Table 3. Plasma total protein, albumin, globulins, AST, and ALT of Californian rabbit does responses to natural additive treatments.

Treatment groups	TP (g/dl)	Alb (g/dl)	Glo (g/dl)	AST (U/l)	ALT (U/l)
Control (C)	4.40 ^c	3.26 ^b	1.13 ^c	11.33	26.33 ^a
Honeybee (HB)	5.91 ^b	3.95 ^a	1.96 ^a	15.500	18.00 ^{ab}
Date palm pollen (DPP)	5.80 ^{bc}	4.20 ^a	1.60 ^b	10.67	22.67 ^{ab}
Bee pollen (BP)	5.80 ^{bc}	4.24 ^a	1.57 ^b	13.00	21.33 ^{ab}
Synbiotic (SY)	6.20 ^a	4.33 ^a	1.87 ^a	8.68	13.68 ^b
Mixture (MIX)	4.77 ^{bc}	3.44 ^b	1.33 ^b	10,00	18.33 ^{ab}
MSE	0.206	0.166	0.098	3.638	4.445
P-value	0.0001	0.0001	0.0001	0.8052	0.0208

a,b, c. Values with different superscripts in the same column, differ significantly ($P < 0.05$). ALT: Alanine aminotransferase, AST: Aspartate aminotransferase, TP: Total protein, Alb: Albumen, Glo: Globulin.

Digestive enzymes

As shown in Table 4 exhibits the activity of amylase and lipase. All bioactive compounds enhanced ($P < 0.05$) the activity of both enzymes. Several studies investigated the effect of probiotic, prebiotic, and bee pollen on the digestive enzymes activity and concluded higher activity than control. **Yang et al (2005)** mentioned that intake of probiotics and prebiotics as a synbiotic mixture by rats significantly improved the ecosystem of the intestinal tract by increasing the activity of digestive enzymes; lipase, lactase, sucrase, and iso-maltase. The higher figures of most metabolites in the current study could be attributed to the elevated activities of digestive enzymes which improve nutrient availability to the animal (**Wang and Gu, 2010**).

Besides, provision of rabbits with bee pollen increased activity of lipase, amylase and protease than controls (**Zeedan et al, 2017**). Concurrently, **Taghian et al (2017)** described the beneficial role of date palm pollen to its higher contents of amino acids and/or its antibacterial activity. Altogether, these actions led to increased nutrient digestibility and growth rate (**El-Speiy et al, 2024**).

Table 4. Blood plasma lipid fractions and digestive Enzymes of Californian rabbit does as affected by natural additive treatments

Treatment groups	TC (mg/dl)	HDL (mg/dl)	LDL (mg/dl)	TG (mg/dl)	Amylase (U/l)	Lipase (U/l)
Control (C)	46.00 ^a	14.57 ^c	7.9 ^a	117.67 ^a	130.66 ^c	140.25 ^c
Honeybee (HB)	37.5 ^b	26.01 ^b	3.49 ^b	40.00 ^c	149.05 ^b	158.40 ^{bc}
Date pollen (DPP)	44.00 ^{ab}	27.9 ^a	4.68 ^b	57.00 ^{bc}	172.4 ^b	184.50 ^b
Bee pollen (BP)	45.67 ^{ab}	28.87 ^b	3.67 ^b	65.67 ^{bc}	169.00 ^b	175.25 ^b
Synbiotic (SY)	43.25 ^{ab}	27.13 ^a	3.25 ^b	64.33 ^{bc}	164.5 ^b	224.30 ^a
Mixture (MIX)	42.33 ^{ab}	25.80 ^b	3.73 ^b	67.33 ^{bc}	149.95 ^b	179.90 ^b
MSE	2.819	0.888	2.902	20.935	21.539	14.362
P-value	0.0043	0.0001	0.002	0.0001	0.0001	0.0001

a, b, c. Values with different superscripts in the same column differ significantly (P<0.05).

TG: Triglycerides, TC: Total cholesterol, HDL: High-density lipoprotein, LDL: low-density lipoprotein, TG: triglycerides.

Reproductive hormones responses to natural additive treatments

As shown in Table 5, revealed that all treatments significantly (P<0.05) elevated E17 β , P4, FSH, LH, and E2/P4 ratio. Contrarily, all treatments reduced (P<0.05) prolactin levels, where synbiotic and mixture resulted in the lowest prolactin levels, followed by honeybee. The improvement of the E2 secretion in does that were given bee derivatives (i.e., honeybee, and bee pollen) ranges between 61-74% over the control values. Moreover, the increase of progesterone ranges between 35-61% over control, with a highest value resulting from bee pollen. In accordance with this finding, **El-Komy *et al* (2021)** documented 15% E2 increase when rabbit does were provided with bee venom.

Many research studies on the effect of honeybee and other bee products (i.e., royal jelly, bee pollen, bee venom) on sex steroid hormones have been published, attributing its effect to the high content of quercetin, caffeic acid, and phenolic compounds in these products (**Zaid *et al*, 2010; Kolesarova *et al*, 2011; Mosavat *et al*, 2014 and Banihani, 2019**). Gonadotrophin hormones (FSH and LH) were also improved remarkably as a function of bioactive compound treatments. As shown in Table 5, the ranges of increase over control for FSH and LH were 49-71%, and 20-33%, respectively. The highest increases of both hormones were evidenced in case of the mixture provision. The augmented effect of the mixture of such bioactive compounds could probably attribute to the facilitative roles of the components on the level of pituitary cells (**Karmali *et al*, 2018**). Likewise, a significant increase of FSH

was documented in women supplemented with probiotic (**Szydłowska et al, 2021**). Furthermore, **Rahman et al (2023)** concluded that provision of probiotic to polycystic ovarian women enormously increased FSH.

Moreover, **Tamemy and Amen (2019)** stated that the provision of rabbit females with date palm pollen appreciably elevated both FSH and LH. They attributed this enhancing effect to the flavonoids that play vital antioxidant and synthesis roles in the mammalian cells. Due to the richness of date palm pollen in phytoestrogens, sterols, carotenoids, and flavonoids, this confers a reproductive health action to this product (**Shahin, 2014**). Seemingly, in rabbits, as in rats and mice, GnRH surge only occurs when estrogen peak and the circadian rhythm clock signals coincide leading to LH surge (**Chappell, 2005**). These together are supported by the concurrent improvement in fertility and the higher litter sizes obtained in the current study in does that were offered bioactive-fortified diets.

Concerning honeybees (HB), **Hashem et al (2021)** reported that honeybees exhibit varieties of nutraceutical compounds (i.e., flavonoids and phenolics) that influence the reproduction functions by encouraging gonadal steroidogenesis, resulting in higher secretion of E2 and P4 in females, and higher androgens in the males. This action on gonads is most probably effective to elicit the gonadotrophin (FSH and LH) secretion from anterior pituitary. Besides, other research attempts have shown great impact of bee pollen (BP) on the ovarian function and local sex steroid hormones secretion (**Kolesarova et al, 2011; Kolesarova et al, 2013 and Attia et al, 2015**). Similar finding to the current result with respect to the reduction of prolactin as a response to the supplementation with bee products was reported previously (**Aoki et al, 2012**).

Likewise, date palm pollen was previously found to reduce levels of prolactin (**Al-Samarrai et al, 2017**). Besides, **Jalalvand et al (2021)** documented a significant reduction of prolactin in women supplied with probiotics bacteria (*Elaeagnus angustifolia* L.). Thus, evidently such bioactive compounds mainly directed their actions on the doe's reproductive traits (**El-Speiy et al, 2024**).

Table 5. Plasma reproductive hormones of Californian rabbit does affected by natural additive treatments

Treatment groups	E17 β (ng/ml)	P ₄ (ng/ml)	E2/P4 Ratio	FSH (ng/ml)	LH (ng/ml)	Prol (ng/ml)
Control(C)	0.166 ^b	0.410 ^c	0.405 ^c	55.26 ^c	46.22 ^c	0.355 ^a
Honeybee (HB)	0.268 ^a	0.554 ^{ab}	0.484 ^b	82.28 ^b	55.38 ^b	0.140 ^c
Date pollen (DPP)	0.293 ^a	0.625 ^a	0.469 ^b	88.79 ^b	58.39 ^a	0.290 ^b
Bee pollen (BP)	0.289 ^a	0.660 ^a	0.439 ^b	89.51 ^b	59.96 ^a	0.295 ^b
Synbiotic (SY)	0.274 ^a	0.610 ^a	0.449 ^b	86.52 ^b	58.45 ^a	0.108 ^c
Mixture (MIX)	0.297 ^a	0.562 ^{ab}	0.528 ^a	94.45 ^a	61.27 ^a	0.109 ^c
MSE	0.002	0.004	0.006	0.009	0.002	0.0954
P-value	0.0001	0.0001	0.0001	0.0001	0.0001	0.0056

a, b, c. Values with different superscripts in the same column differ significantly (P<0.05). E17 β : Estradiol 17 β ; P₄: Progesterone; LH: luteinizing hormone; FSH: follicle stimulating Hormone; Prol: Prolactin.

In conclusion, to improve the productive and reproductive traits of a rabbits raised in developing countries, where lack of feedstuffs prevails, the animal owners must utilize the available bioactive compounds as diet additives. This approach would economically benefit the rabbit raisers and sustain the business for animal protein provision. Also, a multidisciplinary study must be focused to investigate the role (s) of each active constituent of such compounds on the productivity of domestic animals.

REFERENCES

- Abbas, F.A. and Ateya, A. M. (2011).** Estradiol, esteriol, estrone and novel flavonoids from date palm pollen. *Australian Journal of Basic and Applied Sciences* 5(8): 606-614.
- Abd El-Aziz, A., Abo Ghanima, M., Mota-Rojas, D., Sherasiya, A., Ciani, F. and El-Sabrout, K. (2023).** Bee products for poultry and rabbits: Current challenges and perspectives. *Animals*. 13:3517. [https://doi.org/ 10.3390/ani13223517](https://doi.org/10.3390/ani13223517)
- Abdelhady, H. and El-Abasy, M.A. (2015).** Effect of prebiotic and probiotic on growth, immuno-hematological responses and biochemical parameters of infected rabbits with *pasteurell amultocida*. *Benha Veterinary Medical Journal* 28(2): 40- 51. <https://10.21608/BVMJ.2015.31859>
- Abdelnour, S.A., Abd El-Hack, M.E., Alagawany, M., Farag, M.R. and Elnesr S.S. (2019).** Beneficial impacts of bee pollen in animal production, reproduction and health. *J. Anim. Physiol. Anim. Nutr.* 103:477-484. <https://doi:10.1111/jpn.13049>.

- Abdel-Raheem, S. M., Abd-Allah, S. M. S., and Hassanein, K. M. A. (2012).** The effects of prebiotic, probiotic and synbiotic supplementation on intestinal microbial ecology and histomorphology of broiler chickens. *IJAVMS* 6 (4): 277-289. <https://doi.org/10.5455/IJAVMS.156>
- Abdel-Shaheed, M.M., Abdalla, E.S., Khalil1, A.F., and El-Hadidy, E.M. (2021).** Effect of Egyptian date palm pollen (*Phoenix dactylifera* L.) and its hydroethanolic extracts on serum glucose and lipid profiles in induced diabetic rats. *Food and Nutrition Sciences* 12: 147-161. <https://doi.org/10.4236/fns.2021.122013>
- Al-Samarrai, A.H., Al-Samarrai, R.R.H., and Al-Samarrai, N.S.H. (2017).** Biochemical and experimental study for the effect of date palm pollen grains and isolated flavonoids on the levels of steroid hormones in sera of adult female rabbits. *Tikrit Journal of Pure Science* 22(3): 106-110. <https://doi.org/10.25130/tjps.v22i3.719>
- Aoki, M., Wakaoka, Y., Hayashi, H., Kuze, B., Mizuta, K. and Ito Y. (2012).** Effect of lyophilized powder made from enzymolyzed honeybee larvae on tinnitus-related symptoms, hearing levels, and hypothalamus-pituitary-adrenal axis-related hormones. *Ear Hear.* 33(3): 430-436. <https://doi.org/10.1097/AUD.0b013e318234f024>
- Andrade, P., Ferreres, F., and Amaral, M.T. (1997).** Analysis of honey phenolic acids by HPLC, its application to honey botanical characterization. *J. Liq. Chromatogr. Relat. Technol.*, 20: 2281-8. <https://doi.org/10.1080/10826079708006563>
- Attia, Y.A., Bovera, F., El-Tahawy, W.S., El-Hanoun, A.M., Al-Harathi, M.A. and Habiba, H. (2015).** Productive and reproductive performance of rabbits does as affected by bee pollen and/or propolis, inulin and/or mannan-oligosaccharides. *World Rabbit Sci.* 23: 273-282. <https://doi.org/10.4995/wrs.2015.3644>
- Attia, Y. A., Al-Hanoun, A., El-Din, A. E., Bovera, F., Shewika, Y. E. (2011).** Effect of bee pollen levels on productive, reproductive and blood traits of NZW rabbits. *Journal of Animal Physiology and Animal Nutrition* 95(3): 294-303. <https://doi.org/10.1111/j.1439-0396.2010.01054.x>
- Baliga, M.S., Baliga, B.R.V., Kandathil, SM., Bhat, H.P. and Vayalil, P. K. (2011).** A review of the chemistry and pharmacology of the date fruits (*Phoenix dactylifera* L.). *Food Res. Int.* 44(7): 1812-1822. <https://doi.org/10.1016/j.foodres.2010.07.004>

- Banihani, S.A. (2019).** Testosterone in males as enhanced by onion (*Allium Cepa* L.). *Biomolecules* 9(2): 75. <https://doi:10.3390/biom9020075>.
- Biernasiak, J., Śliżewska, K. and Libudzisz, Z. (2010).** Negatywne Skutki Stosowania Antybiotyków. *Postęp Nauk Rol.* 3:105-117.[https:// www.salvum.pl/Stefa-wiedzy/Najczestsze-skutki-uboczne-antybiotykoterapii](https://www.salvum.pl/Stefa-wiedzy/Najczestsze-skutki-uboczne-antybiotykoterapii)
- Birtcher, K.K. and Ballantyne, C.M. (2004).** Measurement of cholesterol: A patient perspective. *Circulation* 110(11): 296-297. [https://doi.org/ 10.1161/01.CIR.0000141564.89465.4E](https://doi.org/10.1161/01.CIR.0000141564.89465.4E)
- Carpes, S. T., Cabral, I. S. R., Luz, C. F. P., Capeletti, J. P., Alencar, S. M. and Masson, M. L. (2009).** Palynological and physicochemical characterizations of *Apis mellifera* L. bee pollen in the Southern region of Brazil. *Journal of Food, Agriculture and Environment* 7(3-4): 667-673.[http://world-food.net/download/journals/2009-issue_3_4/4\(3\).pdf](http://world-food.net/download/journals/2009-issue_3_4/4(3).pdf)
- Cencic, A. and Chingwaru, W. (2010).** The role of functional foods, nutraceuticals, and food supplements in intestinal health. *Nutrients* 2(6):611-625.<https://doi:10.3390/nu2060611>
- Chappell, P.E. (2005).** Clocks and the black box: circadian influences on gonadotropin-releasing hormone secretion. *J.Neuroendocrinol.*17:119-30. <https://doi:10.1111/j.1365-2826.2005.01270.x>.
- De Marco, M., Lopez Salcedo, W., Pastorelli, G., Rossi, R., Corino, C., Bergagna, S., Mellia, E., Gennero, M.S., Biasibetti, E., Capucchio, M.T., Nurisso, S., Tarantola, M., Forneris, G. and Schiavone, A. (2015).** Effects of verbascoside supplemented diets on growth performance, blood traits, meat quality, lipid oxidation and histological features in broiler chickens. *Ital. J. Anim. Sci.* 14:172-178.[https://doi.org/ 10.4081/ijas.2015.3712](https://doi.org/10.4081/ijas.2015.3712)
- Denisow, B. and Denisow-Pietrzyk, M. (2016).** Biological and therapeutic properties of bee pollen. A review. *J. Sci. Food Agric.*96: 4303-4309.<https://doi.10.1002/jsfa.7729>
- Doumas, B.T., Watson, W.A. and Biggs, H.G. (1971).** Albumin standards and the measurement of serum albumin with bromocresol green. *Clin. Chim. Acta.* 31(1):87-96. [https://doi:10.1016/0009-8981\(71\)90365-2](https://doi:10.1016/0009-8981(71)90365-2).
- Duncan, David B. (1955).** Multiple range and multiple F tests." *biometrics*11.1: 1-42. <http://about.jstor.org/terms>
- El-Hammady H. Y., Abuoghaba, A. A., Abd El-Fattah, M.G., and Abd El-Rahman, H.A. (2017).** Semen physical characteristics, blood parameters and some physiological estimates of rabbit bucks administered with bee pollen under Upper Egypt climatic conditions. *Egyptian Journal of Rabbit Science* ,27 (1): 43-64. <https://doi.10.21608/EJRS.2017.41826>

- El-Hanoun, A.M., Kamel, K.I., El-Sepeiy, M.S., and Gad, H.A.M. (2007).** Effect of Egyptian propolis supplementation on productive, reproductive performance and some hematobiochemical parameters and steroid hormones of female rabbits during winter and summer seasons. *In Proc.:4th World Poultry Conference, 27-30 March, 2007. Sharm El-Sheikh, Egypt.*
- El-Komy, A. E., El-Hanoun, A., Abdella, M., El-Sabrou, K. (2021).** Improving the reproductive, immunity and health status of rabbit does using honeybee venom. *J Anim. Physiol. Anim. Nutr.* 105:975-983. <https://doi:10.1111/jpn.13552>
- Ellithy, M., Abdel Fattah, M., and Marwan, A.A. (2022).** Influence of prebiotic, probiotic and synbiotic supplementation on digestibility, hematobiochemical profile and productive performance in Barki lambs. *Egyptian Journal of Nutrition and Feeds* 25(2):199-210. <https://doi:10.21608/ejnf.2022.256903>
- El Rabey, H.A., Al-Seeni, M.N. and Al-Solamy, S.M. (2013).** Bees' honey protects the liver of male rats against melamine toxicity. *Biomed Res Int.* 2013:786051. <https://doi:10.1155/2013/786051>.
- El-Speiy, M.E., Zeitoun, M.M., El-Sawy, M.A., Sadaka, T.A., Abou-Shehema, B.M., Abdella, M.M., Shahba, H.A. and Habib, M.R. (2024).** Bioactive compounds enrichment in rabbit doe's diet pre-and during pregnancy improves productive and reproductive performance and cost-effectiveness under hot climates. *Tropical Animal Health and Production* 54 (7): 170-181.
- Fallahi, S., Rajaei, M., Malekzadeh, K., and Kalantar, S.M. (2015).** Would *Phoenix dactyflera* pollen (palm seed) be considered as a treatment agent against Males' infertility? A systematic review. *Electron Physician.* 7(8): 1590-1596. <https://doi:10.19082/1590>
- Farag, S.A. and El-Rayes, T. K. (2016).** Effect of bee-pollen supplementation on performance, carcass traits and blood parameters of broiler chickens. *Asian J. Anim. Vet. Adv.* 11:168-177. <https://doi:10.3923/ajava.2016.168.177>
- Fossati, P. and Prencipe, L. (1982).** Serum triglycerides determined colorimetrically with an enzyme that produces hydrogen peroxide. *Clin Chem.* 28(10): 2077-2080. <https://pubmed.ncbi.nlm.nih.gov/6812986/>

- Fukushima, M., and Nakano, M. (1995).** The effect of probiotic on fecal and liver lipid classes in rats. *British Journal of Nutrition* 73(1): 701-710. <https://doi:10.1079/bjn19950074>.
- Gawel, S., Wardas, M., Niedworok, E. and Wardas, P. (2004).** Dialdehydmalonowy (MDA) jakows kaźnik procesówperoxydacji lipidów w organizmie [Malondialdehyde (MDA) as a lipid peroxidation marker]. *Wiad Lek.* 57(9-10): 453-455. <https://pubmed.ncbi.nlm.nih.gov/15765761/>
- Gibson, R. G. and Roberfroid, M. B. (1995).** Dietary modulation of the human colonic microbiota: introducing the concept of prebiotics. *J. Appl. Bacteriol.* 125(6): 1401-1412. <https://doi:10.1093/jn/125.6.1401>
- Hannan, A., Barkaat, M., Usman, M., Gilani, W.A. and Sami, W. (2009).** In vitro antibacterial activity of honey against clinical isolates of multi-drug resistant typhoidal salmonellae. *Pakistan J Zool.* 41(1): 1-6. [https://www.zsp.com.pk/pdf/1-6%20\(1\).pdf](https://www.zsp.com.pk/pdf/1-6%20(1).pdf)
- Hashem, N. M., Hassanein, E. M. and Simal-Gandara, J. (۲۰۲۱).** Improving reproductive performance and health of mammals using honeybee products. *Antioxidants (Basel)* 3(10):336-347. <https://doi:10.3390/antiox10030336>
- Hosseini, S.A., Meimandipour, A., Alami, F., Mahdavi, A., Mohiti-Asli, M., Lotfollahian, H. and Cross, D. (2013).** Effects of ground thyme and probiotic supplements in diets on broiler performance, blood biochemistry and immunological response to sheep red blood cells. *Ital. J. Anim. Sci.* 12: 116-120. <https://doi.org/10.4081/ijas.2013.e19>
- Jäger, R., Purpura, M., Farmer, S., Cash, H.A. and Keller, D. (2018).** Probiotic *Bacillus coagulans* GBI-30, 6086 improves protein absorption and utilization. *Probiotics Antimicrob Proteins.* 10(4): 611-615.
- Jalalvand, F., Rezaei, A., Badehnoosh, B., Yaseri, M., Qorbani, M., Emaminia, F. and Shabani, M. (2021).** The Effects of *Elaeagnus angustifolia* L. on the thyroid-stimulating hormone, dehydroepiandrosterone-sulfate, prolactin and cortisol levels in post-menopausal women: A double-blind, randomized, and placebo-controlled study. *Front. Pharmacol.* 12: 654459. <https://doi:10.3389/fphar.2021.654459>
- Karmali, M., Eghbalpour, S., Rajabi, S., Jamilian, M., Bahmani, F., Tajabadi-Ebrahimi, M., Keneshlou, F., Mirhashemi, S.M., Chamani, M., Gelougerdi S.H. and Asemi, Z. (2018).** Effects of probiotic supplementation on hormonal profiles, biomarkers of inflammation and oxidative stress in women with polycystic ovary syndrome: A randomized, double-blind, placebo-controlled trial. *Arch. Iran Med.* 21(1): 1-7

- Kim, M.C. and Kim, M.G. (2023).** Mechanism of blood cholesterol lowering action of paraprobiotics. *Food Suppl. Biomater. Health* 3(1): 1-13. <https://doi.org/10.52361/fsbh.2023.3.e1>
- Kolesarova, A., Capcarova, M., Bakova, Z., Galik, B., Juracek, M., Simko, M. and Sirotkin, A.V. (2011).** The effect of bee pollen on secretion activity, markers of proliferation and apoptosis of porcine ovarian granulosa cells in vitro. *J Environ Sci Health B*. 46(3): 207-212. <https://doi.org/10.1080/03601234.2011.540202>.
- Kolesarova, A., Bakova, Z., Capcarova, M., Galik, B., Juracek, M., Simko, M., Toman, R. and Sirotkin, A.V. (2013).** Consumption of bee pollen affects rat ovarian functions. *J. Anim. Physiol. Anim. Nutr.* 97: 1059-1065. <https://doi.org/10.1111/jpn.12013>.
- Lopez, J. and Alvariño, J.M.R. (2000).** Effects of added caffeine on results following artificial insemination with fresh and refrigerated rabbit semen. *Animal Reproduction Science* 58 (1-2): 147-154. [https://doi.org/10.1016/s0378-4320\(99\)00084-6](https://doi.org/10.1016/s0378-4320(99)00084-6).
- Molan, P.C. and Russell, K. M. (1988).** Non-peroxide antibacterial activity in some New Zealand honeys. *J. Apicult. Res.* 27(10): 62-67. <https://doi.org/10.1080/00218839.1988.11100783>
- Morais, M., Moreira, L., Feás, X. and Estevinho, L.M. (2011).** Honeybee-collected pollen from five Portuguese natural parks: Palynological origin, phenolic content, antioxidant properties and antimicrobial activity. *Food Chem. Toxicol.* 49: 1096-1101. <https://doi.org/10.1016/j.fct.2011.01.020>
- Mosavat, M., Ooi, F.K. and Mohamed, M. (2014).** Stress hormone and reproductive system in response to honey supplementation combined with different jumping exercise intensities in female rats. *BioMed Res. Int.* 2014: 1-6. <https://doi.org/10.1155/2014/123640>
- NRC (1977).** *Nutrient Requirements Of Rabbits*. National Research Council, Washington DC.
- Rahman, N., Hidayat, S.T., Adi Pramono, B., Soerjo Hadijono, R., Oerip Poerwoko, A. and Anityo Mochtar, A. (2023).** Effect of probiotic supplementation on FSH, LH levels and folliculogenesis. *Journal of Biomedicine and Translational Research* 9(1): 19-24. <https://doi.org/10.14710/jbtr.v9i1.16824>

- Ratcliff, W.A., Carter, G.D., Dowsett, M., Hillier, S.G., Middle, J.G., and Reed, M.J. (1988).** Estradiol assays: applications and guidelines for the provision of clinical biochemistry service. *Ann. Clin. Biochem.* 25(5): 466-483. <https://doi.org/10.1177/000456328802500502>
- Reitman, S. and Frankel, S.A. (1957).** Colorimetric method for the determination of serum glutamic oxaloacetic and glutamic pyruvic transaminases. *Am. J. Clin. Pathol.* 28(1): 56-63. <https://doi: 10. 1093/ajcp/28.1.56>.
- S'aric', A., Balog, T., Soboc'anec, S., Kus'ic, B., S'verko, V., Rusak, G., Likic, S., Bubalo, D., Pinto, B., Reali, D. and Marotti, T. (2009).** Antioxidant effects of flavonoid from Croatian *Cystusincanus* L. rich bee pollen. *Food and Chemical Toxicology* 47: 547-554.
- SAS (2002).** SAS user's guide: Statistics version nine. SAS Institutes. Inc. Cary. NC, USA.
- Shahin, F. (2014).** Utilization of date palm pollen as natural source for producing function bakery product. *Egyptian Journal of Agricultural Research* 92: 1457-1470. https://ejar. journals.ekb.eg/article_157004_9a45d4e0c0268a63049a3b8f0476f8bc.pdf
- Simersky, R., Swaczynova, J., Morris, D.A., Franek, M. and Strand, M. (2007).** Development of an ELISA-based kit for the on-farm determination of progesterone in milk. *Vet Med.* 52: 19-28.
- Song, J.G., Baral, K.C., Kim, G.L., Park, J.W., Seo, S.H. Kim, D.H., Jung, D.H., Ifekpolugo, N.L. and Han, H.K. (2023).** Quantitative analysis of therapeutic proteins in biological fluids: recent advancement in analytical techniques. *Drug Deliv.* 30(1): 2183816. <https://doi:10.1080/10717544.2023.2183816>.
- Steel, J.D. and Torrie, J.H. (1984).** Principals and procedures of statistics. A *Biometrical Approach, 4th Edition.* McGraw Hill, USA.
- Sumalatha, K., Saravana, K. A. and Mohana, L. S. (2010).** Review on natural aphrodisiac potentials to treat sexual dysfunction. *Int J Pharm Ther.*, 1: 10-18. https://www.ijptjournal.com/File_Folder/6-14.pdf
- Szydłowska, I., Marciniak, A., Brodowska, A., Loj, B., Cieciewicz, S., Skonieczna-Żydecka, K., Palma, J., Łoniewski, I. and Stachowska, E. (2021).** Effects of probiotics supplementation on the hormone and body mass index in perimenopausal and postmenopausal women using the standardized diet. A 5-week double-blind, placebo-controlled, and randomized clinical study. *Eur. Rev. Med. Pharmacol. Sci.* 25(10):3859-3867.

- Taghian, R.A.S., Abd El-Ati, M.N., Allam, F.M., and Mahmoud, G.B. (2017).** Effect of date palm pollen and bee pollen as growth promoters on the performance of Saidi rams. *Assiut J. Agric. Sci.* 48(5): 86-98. http://www.aun.edu.eg/faculty_agriculture
- Tamemy, S. M. A. L. and Amen, S. H. (2019).** Effect of using different levels of date palm pollen on some hematological and biochemical parameters in local rabbit males. *Plant Archives* 19 (2): 3701-3704. <https://www.cabidigitallibrary.org/doi/pdf/10.5555/20219950736>
- Tuksitha, L., Chen, Y. L.S., Chen, Y. L., Wong, K. Y. and Peng, C. C. (2018).** Antioxidant and antibacterial capacity of stingless bee honey from Borneo (Sarawak). *J. Asia Pac. Entomol.* 21: 563-570. <https://doi.org/10.1016/j.aspen.2018.03.007>
- Wang, Y. and GU, Q. (2010).** Effect of probiotic on growth performance and digestive enzyme activity of Arbor Acres broilers. *Res. Vet. Sci.* 89: 163-167. <https://doi.org/10.1016/j.rvsc.2010.03.009>
- White, J.W. (1975).** Composition of Honey. In: Crane E, Ed. Honey, and a comprehensive survey. London: *Bee Research Association and Chalfont St Peter*;: 157-206. <https://www.scirp.org/reference/references/papers?Referenceid=681687>
- Yang, S.C., Chen, J.Y., Shang, H. F., Cheng, T.Y., Tsou, S. C. and Chen, J.R. (2005).** Effect of synbiotics on intestinal microflora and digestive enzyme activities in rats. *World Journal of Gastroenterology*, 11(47): 7413-7417. <https://doi:10.3748/wjg.v11.i47.7413>
- Zaid, S.S.M., Sulaiman, S.A., Sirajudeen, K.N.M. and Othman, N.H. (2010).** The effects of Tualang honey on female reproductive organs, tibia bone and hormonal profile in ovariectomized rats: Animal model for menopause. *BMC Complement. Altern. Med.* 10: 82. <https://doi:10.1186/1472-6882-10-82>
- Zeedan, K., Ali, N.G.E.M., Abuoghaba, A.A. and El-Kholy, K. (2017).** Effect of bee pollen at different levels as natural additives on immunity and productive performance in rabbit males. *Egypt. Poult. Sci.* 37(1): 213-231. https://epsj.journals.ekb.eg/article_6778.html
- Zoli, A., Lizzio, M.M., Ferlisi, E.M., Massafra, V., Mirone, L., Barini, A., Scuderi, F., Bartolozzi, F. and Magaró, M. (2002).** ACTH, cortisol and prolactin in active rheumatoid arthritis. *Clin. Rheumatol.* 21(4): 289-293. <https://doi:10.1007/s100670200076>

تعزيز المناعة وحالة الأكسدة والتمثيل الغذائي ومستوى هرمونات التناسل بأضافة بعض الاضاقات الطبيعية لعليقة إناث الأرناب

محمد السيد السبيعي^١ - محمد عبد العزيز الصاوي^١ - طارق أمين صدقه^١ - بهاء محمد السيد أبو شحيمه^١ - حسام عبد المنعم شهبه^١ - محمود رشاد حبيب^١ - مصطفى محمد عبد اللاه^١ - أحمد محمد تمام^٢ - مصطفى محمد زيتون*^٣

- ١- قسم بحوث الأرناب - معهد بحوث الأنتاج الحيواني - مركز البحوث الزراعية - مصر.
- ٢- قسم تغذية الدواجن - كلية الزراعة - جامعة عين شمس - مصر.
- ٣- قسم الإنتاج الحيواني والسمكي - كلية الزراعة بالشاطبي - جامعة الإسكندرية - مصر.

يهدف البحث الى إيجاد مواد طبيعية محلية تعزز المناعة وحالة الأكسدة والتناسل في إناث الأرناب الكاليفورنيا. تم توزيع ٧٢ أنثى بالغة على ٦ معاملات تجريبية عشوائيا (١٢ أنثى بكل معاملة). كل الإناث غُذيت على العليقة الأساسية كما تم تجريعها فمويا ٣ مل من الماء المقطر (تحتوى على الإضافة) قبل التلقيح ب ١٠ أيام وحتى اليوم الـ ٢٨ بعد التلقيح بشكل مستمر. مجموعة المقارنة (الكونترول) لم تتلقى أى إضافات، بينما تم تجريع المجموع الأخرى من الثانية حتى السادسة الكميات الأتية بالترتيب: ٥٠ مجم عسل نحل، ٥٠ مجم طلع النخل، ٢٠٠ مجم حبوب لقاح النحل، ٣ مل سينبيوتيك، أو خليط من كل المواد السابقة. وقد أوضحت النتائج أن جميع المعاملات أدت إلى تحسن معنوى بمستوى ٥% للمناعة وحالة الأكسدة، كما لوحظت زيادة معنوية بمستوى ٥% فى محتوى بلازما الدم من كل من البروتين الكلى والألبومين وكذلك الجلوبيولين، بينما إنخفضت مستويات إنزيمات الكبد، أيضا أدت جميع المعاملات لإنخفاض الكوليسترول منخفض الكثافة (الضار) والدهون الثلاثية بينما زاد مستوى الكوليسترول عالى الكثافة مقارنة بالكنترول. كما أدت المعاملات إلى زيادة مستوى هرمونات التناسل المفرزة من المبيض (البروجستيرون والاستراديول) وتلك المفرزة من النخامية (FSH, LH) كما أدت جميع المعاملات إلى خفض مستوى هرمون البرولاكتين مقارنة بالكنترول. **الاستنتاج:** إضافة كل من منتجات النحل أو طلع نخيل البلح أو السينبيوتيك أو الخليط في مياه شرب إناث الأرناب أدت لتحسين المناعة والصفات الإنتاجية.

التوصية: نظراً لنقص المواد العلفية في الدول النامية، بالإضافة إلى قسوة المناخ، فإن هذه العوامل مجتمعة تشكل تحدياً كبيراً أمام مربى الأرناب للحصول على الإنتاج الاقتصادي. علاوة على ذلك، وكما هو معروف فإن الفراء الكثيف الموجود على جسم الأرناب يشكل عبئاً أكبر على إنتاجيته. ومن أجل تعزيز الصفات الإنتاجية والإنجابية للأرناب، يجب على مربى الأرناب استخدام المركبات النشطة بيولوجيا المتوفرة كإضافات غذائية، وخاصة تلك التي استخدمها في الدراسة. وهذا النهج من شأنه أن يفيد مربى الأرناب اقتصادياً ويدعم زيادة انتاج البروتين الحيواني. يجب أن تركز على دراسات متعددة التخصصات لتحديد دور (أدوار) كل مكون نشط من هذه المركبات على الإنتاجية الحيوانية.

الكلمات المفتاحية: منتجات نحل العسل، طلع النخل، المناعة، البروبيوتيك، الكفاءة الانتاجية، الأرناب.