

Table (5). Carcass traits of New Zealand White rabbits as affected by addition algae species and levels.

Items	Control	Algae species (g/ Kg diet)				±SE	P<
		<i>Spirulina platensis</i>		<i>Chlorella vulgaris</i>			
		(0.75)	(1.5)	(0.75)	(1.5)		
Slaughter weight at 12-wks (g)	1827.3 ^c	1935.6 ^{ab}	2003.3 ^a	1903.6 ^{bc}	1865.3 ^{bc}	26.5	0.018
Hot carcass weight (g)	1017.3 ^c	1112.3 ^a	1133.0 ^a	1100.3 ^{ab}	1052.3 ^{bc}	16.1	0.029
Liver weight (g)	74.96	68.13	75.46	66.10	57.33	8.23	0.544
Liver weight (%)	4.10	3.51	3.75	3.46	3.06	0.399	0.494
Kidney weight (g)	15.16	14.63	15.53	12.96	17.13	1.02	0.136
Kidney weight (%)	0.83	0.75	0.77	0.68	0.91	0.052	0.084
Heart weight (g)	6.96	9.70	7.93	7.06	7.16	0.93	0.269
Heart weight (%)	0.383	0.506	0.393	0.370	0.383	0.050	0.350
Total giblet weight (g)	90.43	92.46	98.93	86.13	81.63	10.13	0.793
Total giblet weight (%)	4.93	4.77	4.93	4.52	4.36	0.498	0.898
Dressing weight (g)	1107.7 ^c	1204.8 ^{ab}	1231.9a	1186.4 ^{ab}	1133.9 ^{bc}	21.85	0.014
Dressing weight (%)	60.62	62.23	61.50	62.33	60.78	0.695	0.332

^{a, b, c} Means within the same row with different litters differ significantly at P<0.05.