

**Table 1:** Included case of rabbit osteosarcoma with extracted information

Reference	Age	Breed	Wt	Sex	LPT	Metastasis	Symptoms	Diag. Tech	Treatment	Outcome
<b>Gibson and Donnelly, 2011</b>	18 months	NZW	2.3 kg	Fe	Proximal aspect right femur	Lungs	Anorexia Decreased urine output Decreased stool output Right hind limb lameness	Radiography	None	Euthanized
<b>Weiss and Müller, 2011</b>	2 years	Not mentioned	1.2 kg	M.	Spine	None identified	Anogenital soiling Dermatitis. Ataxia of pelvic limbs Swelling dorsal sacral region. Atrophy pelvic limb muscles	Radiography	None	Euthanized
<b>Ishikawa et al., 2012</b>	7 years	Cross bred	2.6 kg	Fe	Tarsal joint region -involving distal tibia	Pulmonary micrometastases None in liver	Bleeding in tarsal region right hind limb Swelling in tarsal region right hindlimb	Radiography	Mass excision (CO2-laser)	Died 3 days post-surgery
<b>Wijesundera et al., 2013</b>	9 years	Mixed	1.8 kg	M	Subcutaneous in left forelimb (extraskkeletal)	None	Hemorrhage & edema around declaw left forelimb (initial). Ulcerating mass at resection site (2 months after resection)	Radiography	Surgical removal (primary tumour). Amputation of limb one month after recurrence	Recurrence 2 months after resection. Alive
<b>Higgins et al., 2015</b>	8 years	miniature Rex	1.7 kg	M	Tibia or fibular head	No evidence of pulmonary metastases at presentation No evidence of metastases 4 months after amputation. No evidence of metastases 6 months after amputation.	Lameness  Atrophy left pelvic limb musculature	Radiography  Fine needle aspiration	Pre-amputation: tramadol, meloxicam, gabapentin Amputation of limb after three days	Active 2 weeks after operation. Died 190 days after surgery recovering from gastrotomy for gastric bloat

<b>Continue Table 1</b>										
<b>Reference</b>	<b>Age</b>	<b>Breed</b>	<b>Wt</b>	<b>Sex</b>	<b>LPT</b>	<b>Metastasis</b>	<b>Symptoms</b>	<b>Diag. Tech</b>	<b>Treatment</b>	<b>Outcome</b>
<b>van Praag, 2015</b>	1 year	Rex	Not mentioned	Fe	Pelvis, lumbar vertebra, tail base	Multiple abdominal, chest and pulmonary metastases	Lameness  Paresis of hind limbs	Radiography  Biopsy	None mentioned	Euthanized
<b>Makishima et al., 2020</b>	8 years, 9 months	Lop-eared	1.9 kg	M	Intraocular, extraskelatal	None (from primary tumor) Separate primary pulmonary neoplasms	Enlargement of left eye Exophthalmos Respiratory failure	None	None mentioned	Died due to respiratory failure (from pulmonary neoplasms)
<b>Miwa et al., 2019</b>	4 years	Mixed	Not mentioned	M	Mandibular gingiva	Pulmonary metastases	Not mentioned	Computed tomography	Partial mandibulectomy (Primary) Hemi-mandibulectomy (Recurrence)	Died 4 months after diagnosis
<b>Miwa et al., 2019</b>	4 years	Lion-headed	Not mentioned	M	Maxillary gingiva	Pulmonary metastases	Not mentioned	Radiography	Laser resection (Primary) Radiation (Recurrence)	Died 45 months after diagnosis
<b>Miwa et al., 2020</b>	7.2 years	Not mentioned	Not mentioned	M	Intranasal	Not mentioned	Nasal discharge	Radiography	Rhinostomy	Unknown

Fe: Female, M: Male, NZW: New Zealand White , Wt: Weight, LPT: Location primary tumour, Diag. Tech : Diagnostic techniques